

## Directions for use

**Application:** Boom Spray, Fertigation or Aerial

Crop	Rate L/Ha	Notes
<b>Broad Acre Grasses</b> Cereals, sorghum, corn, sugar cane.	1.0 – 3.0 1 to 2 Applications	Apply to non-stressed plants from the second internode elongation up to head emergence.
<b>Broad Acre Dicots</b> Cotton, potatoes, clovers, lucerne, peanuts, canola, pulses, sunflowers	1.0 – 3.5 1 to 4 Applications	Apply to non-stressed plants at the commencement of their reproductive phase.
<b>Horticulture</b> Vines, Citrus, Vegetables, Trees – nut & fruit	1.0 – 4.0 1 to 4 Applications	Apply to plants at the commencement of their reproductive phase at a minimum dilution of 1:100 with water.

### Standard mixing & spraying instructions:

1. **Agitate contents well before use.**
2. Fill tank to a minimum of two thirds with water.
3. Add other nutrients as prescribed under agitation.
4. Add BioAg Fruit & Balance® under agitation, bring volume up to full.
5. Before tank mixing, jar test all added nutrients for compatibility using mixing instructions. Check for physical changes.
6. **Use within 48 hours of dilution.**
7. Rinse sprayer after use.

NOTE: The above suggested rates should be used as a guide only. BioAg does not warrant that these rates will necessarily produce the optimal or expected outcome. Each farmer's climatic conditions, before, during and after application, water quality, soil and plant types, application processes and practices may differ and require adjustments to the application rates to ensure optimum results.

Manufactured by BioAg Pty Ltd ABN 58 086 880 211  
22-26 Twynam St Narrandera NSW 2700 Australia  
T +61 2 6958 9911 F +61 2 6959 9922  
E sales@bioag.com.au W www.bioag.com.au

**Keep out of reach of children. Do not swallow.  
The spray form this product may act as an irritant.  
Avoid inhalation and contact with the eyes and skin.**



# Fruit & Balance® Foliar Fruiting Supplement



Australian made  
Australian owned and operated



Produced by aerobic fermentation BioAg Fruit & Balance® has been developed to aid the fruiting and reproductive processes in the plant and to assist in the expansion of the leaf and soil foodwebs.

### Storage and handling:

1. Store all containers out of sunlight as the product is light and temperature sensitive.
2. The product has a shelf life of 18 months at temperatures below 30°C.

### Conditions of sale:

1. The product has a limited shelf life and must be stored in accordance with the storage and handling directions to preserve shelf life. Failure to store correctly or use after the expiry of shelf life is a breach of the conditions of sale. As BioAg cannot exercise control over storage, handling, and the exercise of personal judgment, skill and expertise in the mixing, application or use, or weather, or animal condition before, during or after application, all of which may affect the performance of the product, BioAg, to the maximum extent permitted by law, will not accept any liability for any failure in performance of the product, or any losses or damages whatsoever.
2. Except as expressly provided in these Conditions of Sale, no warranty, condition, undertaking or term, express or implied, statutory or otherwise, as to the quality or fitness for purpose of the goods is given or assumed by BioAg and all such conditions, warranties and liabilities (including liability as to negligence) express or implied relating to the goods are hereby excluded to the extent permitted by law. To the extent that liability can be shown (under law) it is limited to the replacement of goods or their equivalent monetary value only.

**Analysis:** Please contact BioAg for a complete product analysis.

Batch No / DOM:					
Net Contents	10 L	20 L	60 L	200 L	1,000 L